

Cumulant ratios and their scaling functions for Ising systems in a strip geometry

¹Andrzej Drzewiński, ²Jacek Wojtkiewicz

¹*Institute of Low Temperature and Structure Research, Polish Academy of Sciences,
P.O.Box 1410, 50-950 Wrocław 2, Poland.*

²*Department for Mathematical Methods in Physics,
Warsaw University, 00-682 Warszawa, Poland.*

(February 1, 2008)

We calculate the fourth-order cumulant ratio (proposed by Binder) for the two-dimensional Ising model in a strip geometry $L \times \infty$. The Density Matrix Renormalization Group method enables us to consider typical open boundary conditions up to $L = 200$. Universal scaling functions of the cumulant ratio are determined for strips with parallel as well as opposing surface fields.

PACS numbers: 05.50.+q, 68.35.Rh, 75.10.H

Introduction. An *universality principle* is a cornerstone of contemporary theory of phase transitions. According to this principle, the following sorts of quantities are universal: critical exponents, certain amplitude ratios and scaling functions [1]. They differ each from other in their status. The (bulk) critical exponents are independent on boundary conditions, whereas two other groups are dependent.

The critical exponents are known for many models (both exactly and approximately). The collection of results available for amplitude ratios is also rich, but significantly smaller than for exponents; see in Ref. [1] for exhaustive information.

Among amplitude ratios, so called *cumulant ratios* are of great importance. They supply some information on scaling functions (cumulants are proportional to derivatives of these functions at zero values of argument(s)); they measure deviation of magnetization fluctuations at criticality from gaussian distribution; moreover, they are closely related to some versions of renormalization group [2] (it gains also reflection in terminology: cumulant ratios are customarily termed “renormalized coupling constant” in field theory). Cumulant ratios have also been used to locate the critical points and critical lines in many models [2–5].

Most results for cumulants have been obtained for (partially or completely) periodic boundary conditions. For most extensively studied the two-dimensional Ising model, numerous results are available [2,5–13]. However, there are also other, very natural “open” boundary conditions: “free” (no surface fields), “wall++” (infinite parallel surface fields), “wall+-” (infinite opposing surface fields). For these open boundary conditions, the number of results is very small. Only papers we know where such results are available are [2,4]. Motivated by this situation, we state the aim of this paper:

Calculation of universal cumulant ratios for the two-dimensional Ising model in a strip geometry under the following boundary conditions: “free”, “wall++”,

“wall+-”.

We have calculated cumulant ratios using method called the Density Matrix Renormalization Group (DMRG). Since the DMRG is most powerful for open boundary conditions, it is particularly suited for our goals.

Definition of cumulants. We consider the two-dimensional Ising system on a square lattice in a strip geometry (L is width of the strip and N is its length) with the Hamiltonian

$$\mathcal{H} = -J \left[\sum_{\langle i,j \rangle} s_i s_j - H \sum_i s_i - H_1 \sum_i s_i - H_L \sum_i s_i \right], \quad (1)$$

where the first sum runs over all nearest-neighbour pairs of sites while the last two sums run over the first and the L -th column, respectively. H is the bulk magnetic field, whereas H_1 and H_L are the surface fields. H , H_1 and H_L are dimensionless quantities (all of them are measured in units of J). In the course of calculations of the cumulant in the thermodynamic limit, two limiting processes are taken: $T \rightarrow T_c$ and $L \rightarrow \infty$. In general, a value of the cumulant do depend on ordering of these limits [12]. In our paper we analyze so-called “massless” case (analogously as in [12,14]): $T = T_c = 2/\ln(1 + \sqrt{2}) \simeq 2.269185$ followed by $L \rightarrow \infty$. Therefore, we do not notice the temperature dependence below. We also drop (as unnecessary) explicit dependence on surface fields until discussion of scaling functions.

We consider the ratio of moments of magnetization proposed by Binder [2]. Definitions of cumulant ratios for a system in a strip geometry have been widely presented in the literature [2,5,8,11]. Let us first define

$$U_L = \lim_{N \rightarrow \infty} [N(1 - \frac{1}{3} \langle M^4 \rangle / \langle M^2 \rangle^2)], \quad (2)$$

where $M = \sum_i s_i$ is the total (extensive) magnetization. Then, the cumulant ratio A_U in question is

$$A_U = \lim_{L \rightarrow \infty} L^{-1} U_L. \quad (3)$$

An equivalent (but more convenient for us) formula for the above cumulant is as follows: Let $\lambda(L; H)$ be the largest eigenvalue of transfer matrix for the strip of width L ($-T \log \lambda(L; H)$ is the free energy for one column of spins). Define

$$m_2(L) = \frac{d^2}{dH^2} \log \lambda(L; H)|_{H=0}, \quad (4)$$

$$m_4(L) = \frac{d^4}{dH^4} \log \lambda(L; H)|_{H=0}. \quad (5)$$

Then our cumulant is equal to [5]

$$A_U = \lim_{L \rightarrow \infty} r(L) \equiv \lim_{L \rightarrow \infty} m_4(L)/3Lm_2^2(L). \quad (6)$$

Our method of calculation is based directly on the definition (6). We first find logarithms of the largest eigenvalue $\lambda(L; H)$ for some values of H (at fixed L). Next, we calculate numerically derivatives (4) and (5), then the ratio $r(L)$, and finally perform the extrapolation $L \rightarrow \infty$.

Some technical details of calculations. We use the DMRG method for calculations of $\log \lambda(L; H)$. Originally, this method has been proposed by White [15] for finding accurate approximations to the ground state and the low-lying excited states of quantum chains. Its heart is recursive construction of the effective Hamiltonian of a very large system using a truncated basis set, starting from an exact solution for small systems. Later, the DMRG was adapted by Nishino [16] for two-dimensional classical systems. The DMRG has been applied successfully to many different problems and now it can be treated as a standard method, which is very flexible, relatively easy to implement and very precise. For a comprehensive review of a background, achievements and limitations of DMRG, see in Refs. [17,18].

A factor crucial for precision of DMRG is so-called *number of states kept* m , describing the dimensionality of effective transfer matrix [15,17]; the larger number of states kept, the more accurate the value of the free energy. Using $m = 50$ we can calculate the free energy with accuracy of the order 10^{-12} for strips of width of the order $L = 200$. This is an one order more than size of systems available by exact diagonalization of transfer matrix. This fact is crucial for us, because of using the extrapolation procedures. In our calculations we apply the *finite system algorithm*, developed by White for studying finite systems [15]. Additional factor determining the accuracy of the method is *number of sweeps*, i.e. number of iterations made in order to obtain self-consistency of results. Our numerical experience shows that in most cases, it is sufficient to apply only one sweep (although in the "wall+-" case two sweeps are necessary – see below).

In our calculations of cumulant ratios, we have also a factor limiting accuracy that is independent on the DMRG method: *accuracy of numerical differentiation*. In the procedure of numerical differentiation, a suitable choice of increment ΔH of an argument is of crucial importance. It is clear that ΔH should be taken as small as possible; on the other hand, due to finite accuracy in calculation of λ , an error of difference quotient increases with decreasing ΔH . The increments used in our calculations have been determined as a compromise between above two tendencies. Additional factor determining the accuracy of numerical differentiation, is a number of points used to calculate the derivative. We use formulas where a derivative is determined from the second-order Taylor expansion (i.e. we need $n+3$ values of function for n -th derivative; this way, an accuracy is of the order $\mathcal{O}((\Delta H)^3)$.) Therefore, the m_2 was determined from 5 points (3 points in symmetrical case, i.e. $f(H) = f(-H)$) and from 7 points for m_4 (4 points when the symmetry was present).

We have tested correctness of our calculations in several ways. One of them was L -dependence on derivatives m_2 and m_4 . Finite-Size Scaling (FSS) theory [1,19] predicts the following dependence on the n -th derivative of the free energy as a function of the system size L :

$$\frac{d^n f}{dH^n}(L)|_{H=0} \sim L^{-\tilde{d}+n\Delta/\nu}, \quad (7)$$

where we have $\Delta = 15/8$ and $\nu = 1$ for the two-dimensional Ising model. $\tilde{d}$ is a dimension of system in "finite-size direction", i.e. it is a number of linearly independent directions along which a size of the system is finite. For finite systems (for instance a torus), $\tilde{d}$ is equal to space dimension of the system. In our case, the system is infinite in one direction (along the strip) and finite in the second direction (across the strip), so we have to take $\tilde{d} = 1$. This assumption gives the following predictions for derivatives:

$$m_2 \sim L^{\rho_2}, \quad m_4 \sim L^{\rho_4}, \quad (8)$$

where $\rho_2 = 11/4$ and $\rho_4 = 13/2$.

An extrapolation procedure has been performed with use of the powerful BST method [20].

Results: the "free" case. It corresponds to zero surface fields $H_1 = H_L = 0$ in the formula (1). We have performed calculations for L in the range $160 \leq L \leq 200$ with step 10; these values of L were taken in all situations. We took an increment of "bulk" magnetic field $\Delta H = 5 \times 10^{-6}$, $m = 50$ and one sweep. The results are listed in the Table.

As a byproduct, we have tested the FSS predictions for L -dependence of derivatives m_2 and m_4 . Values of corresponding exponents (see Eq. (8)) are: $\rho_2 = 2.7495(3)$ and $\rho_4 = 6.50(3)$, so predictions of FSS are confirmed in excellent manner. The same conclusion is true in next two situations.

As another test of correctness (and quality) of DMRG results, we have calculated ratios by immediate *numerical diagonalization of transfer matrix* for $10 \leq L \leq 18$ (L even; these values of L have also been used in the next cases). We proceeded as above, i.e. by calculation of logarithm of the largest eigenvalue for some values of bulk field H , followed by numerical differentiation of $f(H)$ and computation of ratio and extrapolation, without any “renormalization”. We took an increment $\Delta H = 10^{-4}$. We have obtained $A_U = -1.094(1)$; $\rho_2 = 2.746(1)$; $\rho_4 = 6.46(1)$. It is seen that the results are fully consistent with the DMRG calculations but less precise; we have the same situation for two other boundary conditions.

Results: the “wall++” case. The “wall++” boundary condition corresponds to the assumption that all boundary spins have the same value and sign. It is equivalent to putting $H_1 = H_L = \infty$ in (1). Numerical experience suggests that it is sufficient to take $H_1 = 10$ – for larger values of H_1 the changes of the free energy are negligible [21].

The “wall++” configuration is more intricate, from numerical point of view, than “free” system. The complication is due to the fact that, for parallel surface fields of the same sign, the maximum of the free energy $f(H)$ does not appear for $H = 0$ but it is shifted to a certain non-zero value $H_0(L)$. This phenomenon is called the *capillary condensation* [21,22]. In order to calculate derivatives and ratios at zero magnetization (i.e. at the maximum of the free energy), we first have to find its position $H_0(L)$. FSS predicts [22] the following dependence: $H_0(L) \sim L^{-\Delta/\nu}$. From our DMRG calculations, we have obtained the value $\Delta/\nu = 1.8749(2)$.

For the “wall++” configuration the free energy is not longer a symmetric function of the bulk field H , so we have been forced to calculate m_2 from 5 points and m_4 from 7 points. We have taken an increment $\Delta H = 5 \times 10^{-6}$, $m = 50$ and one sweep. The results are presented in the Table. For exponents of m_2 and m_4 , we have obtained: $\rho_2 = 2.7504(3)$ and $\rho_4 = 6.5024(3)$. The precision of these results is a little bit less than in the “free” case, although still very satisfactory (three significant digits). However, it should be stressed that here we must do much more numerical computations than in the “free” case, so some lack of precision is inevitable.

Exact diagonalization of transfer matrix gave the following values: $A_U = 0.45(4)$; $\rho_2 = 2.75(1)$; $\rho_4 = 6.5(2)$.

Results: the “wall+-” case. One of important physical implications of “+-” boundary condition is the presence of an *interface* between “+” and “-” phases in the system. It causes large fluctuations, which have an implication in numerical practice, namely, *two sweeps* are necessary to ensure self-consistency of results. In our calculations, the value of surface field $H_1 = 100$ an increment $\Delta H = 2 \cdot 10^{-5}$, and $m = 40$ were taken. The results are listed in the Table. Values of exponents are:

$\rho_2 = 2.7502(2)$, $\rho_4 = 6.502(2)$.

The procedure of exact diagonalization of transfer matrix gave the following values: $A_U = -0.305(2)$; $\rho_2 = 2.755(1)$; $\rho_4 = 6.50(2)$.

As a matter of some interest, let us remark that for the “wall+-” boundary condition the L dependence is much weaker than for the “free” and “wall++” situations.

Table. Values of cumulant ratios for some values of L

L	$r(L)$, free	$r(L)$, wall++	$r(L)$, wall+-
160	-1.098525	0.462556	-0.304831
170	-1.098234	0.462225	-0.304859
180	-1.097964	0.461859	-0.304883
190	-1.097723	0.461525	-0.304902
200	-1.097481	0.461133	-0.304915
∞	-1.0932(3)	0.455(2)	-0.3050(1)

Scaling functions for ratios. The “wall++ / +-”-type conditions can be treated as limiting case of the system with equal *finite* parallel/antiparallel (+ / + -) surface fields. Another limiting case is the “free” boundary condition, where the values of surface fields are set to zero. One can expect that for intermediate situations, i.e. finite values of boundary field H_1 , cumulants would be smooth functions of H_1 . Particularly interesting are *scaling properties* of these functions. The scaling theory predicts that at criticality, the system depends only on one variable, namely dimensionless combination $\zeta = LH_1^2$ [23]. In the other words, we expect that the cumulant r should depend on the surface field H_1 and strip width L only by combination ζ . We have calculated $r(L, H_1)$ for both “++” and “+-” boundary conditions for $L = 40, 80, 120$ using $m = 40$, and at full range of scaling variable. The results are presented on Figs. 1a and 1b. It is seen that scaling properties are confirmed in excellent manner.

Limiting values of these functions (i.e. for $\zeta = 0$ and $\zeta \rightarrow \infty$) are fully consistent with our more precise calculations, although convergence of ratio to its limit value is much faster for “wall+-” than for “wall++”. As far we know, scaling functions for cumulants have been almost not studied so far. Only exception is paper [24]; however, the authors consider scaling functions different from ours.

Summary. We have calculated cumulant ratios for Ising strips with three natural boundary conditions, almost not studied so far: “free”, “wall++” and “wall+-” situations. We have applied the Density Matrix Renormalization Group method followed by numerical differentiation and extrapolation $L \rightarrow \infty$. We claim that our results are very precise (three or four significant digits). The precision is comparable with three other “top quality” methods used in similar calculations: Monte Carlo [12], some versions of Renormalization Group [13] and analysis of high-temperature series [10].




FIG. 1. The cumulant ratio as a function of scaling variable $\zeta = LH_1^2$: a) the “++” boundary condition; b) the “+-” boundary condition. Notice much weaker L dependence for “+-” than for “++” situation. Moreover, the convergence of ratio to its limit value, when $\zeta \rightarrow \infty$ is much faster for “+-” than for “++”. For the latter case, the saturation is achieved for $\zeta \approx 500$ (outside the range of plot).

We have also calculated the quantity which apparently escaped the attention so far (at least for boundary conditions considered by us), namely, the scaling functions for cumulants. Such functions give information of how finite surface fields influence values of cumulants. This influence is significant – in one case (“++”) even the sign of cumulant changes upon growth of surface field. We do not know how fundamental this phenomenon is. At first glance, it seems to be related to the lack of symmetry (i.e. $f(H_0 + H) \neq f(H_0 - H)$) and to a non-zero value of the third derivative of the free energy f with respect to H at H_0 . However, the other explanations are not excluded and we will discuss it elsewhere.

Natural lines of continuation of our investigations are: testing of universality of cumulants and scaling functions (for other models in the two-dimensional Ising universality class, for example the hard squares model) and calculation of higher cumulants. This work is currently in progress.

Most of our numerical calculations were performed on Pentium II machines under Linux. We used ARPACK package to calculate eigenvalues.

Acknowledgments. This work was partially funded

- [1] V. Privman, P.C. Hohenberg, and A. Aharony, in *Phase Transitions and Critical Phenomena* edited by C. Domb and J.L. Lebowitz (Academic Press, London 1991), vol. 14, p. 1.
- [2] K. Binder, Z. Phys. B - Condensed Matter **43**, 119 (1981).
- [3] G. Kamieniarz, P. Kozłowski, and R. Dekeyser, Acta Magnetica **XI**, 71 (1996/1997); G. Kamieniarz, P. Kozłowski, and R. Dekeyser, Phys. Rev. E **55**, 3724 (1997).
- [4] K. Binder, D. P. Landau, and A. M. Ferrenberg, Phys. Rev. E **51**, 2823 (1995).
- [5] H. Saleur and B. Derrida, J. Physique **46**, 1043 (1985).
- [6] A.D. Bruce, J. Phys. C **14**, 3667 (1981).
- [7] A.D. Bruce, J. Phys. A **18**, L873 (1985).
- [8] T.W. Burkhardt and B. Derrida, Phys. Rev. B **32**, 7273 (1985).
- [9] R.C. Desai, D.W. Heermann, and K. Binder, J. Stat. Phys. **53**, 795 (1988).
- [10] M.E. Fisher, S.-N. Lai, and S.-Y. Zinn, Phys. Rev. E **54**, 1176 (1996).
- [11] G. Kamieniarz and H. Blöte, J. Phys. A **26**, 201 (1993).
- [12] J. Salas and A. Sokal, cond-mat/9904038, to appear in J. Stat. Phys.
- [13] A.I. Sokolov and E.V. Orlov, Phys. Rev. B **58**, 2395 (1998).
- [14] J.-K. Kim, A. Patrascioiu, Phys. Rev. D **47**, 2588 (1993).
- [15] S.R. White, Phys. Rev. Lett. **69**, 2863 (1992); S.R. White, Phys. Rev. B **48**, 10345 (1993).
- [16] T. Nishino, J. Phys. Soc. Jpn. **74**, 3598 (1995).
- [17] *Lecture Notes in Physics* edited by I. Peschel, X. Wang, M. Kaulke, and K. Hallberg (Springer-Verlag, Berlin 1999), vol. 528.
- [18] K. Hallberg, cond-mat/9910082.
- [19] M.N. Barber, in *Phase Transitions and Critical Phenomena* edited by C. Domb and J. L. Lebowitz (Academic Press, London 1983), vol. 8, p. 145.
- [20] M. Henkel and G. Schütz, J. Phys. A **21**, 2617 (1988).
- [21] E. Carlon, A. Drzewiński, and J. Rogiers, Phys. Rev. B **58**, 5070 (1998).
- [22] M.E. Fisher and H. Nakanishi, J. Chem. Phys. **75**, 5857 (1981); H. Nakanishi and M.E. Fisher, J. Chem. Phys. **78**, 3279 (1983).
- [23] M.E. Fisher and P. de Gennes, C.R. Acad. Sci. Ser. B **287**, 207 (1978).
- [24] Y. Okabe and M. Kikuchi, Int. J. Mod. Phys. C **7**, 287 (1996).